

2006 Florida Youth Tobacco Survey

Bay County

Changes and Trends from 2000 to 2006




The Florida Youth Tobacco Survey (FYTS) is a statewide, school-based confidential survey that has been conducted annually since 1998. The FYTS collects information regarding the statewide prevalence of tobacco use, attitudes, and related behaviors among middle and high school students throughout Florida and tracks those behaviors over time. The FYTS has been administered annually at the state level since 1998 and has been administered at the county level in 2000, 2002, and now 2006. With three years of county level data, we are now able to track changes within each county.

In Bay County, 628 middle school students and 684 high school students in 15 public schools completed the 2006 FYTS. This report presents the county level data in 2000, 2002, and 2006. State and county prevalence from 2000 and 2002 were included for comparison. The county level data and state level data are weighted to represent the entire population of Bay County public middle and high school students and Florida public middle and high school students respectively. These data can be used by county health departments, school districts, and other community organizations and citizens to estimate the magnitude of youth tobacco use in Bay County, as well as to compare tobacco use, attitudes, and related behaviors in Bay County to those for Florida as a whole.

Current Tobacco Use

Current Tobacco Use


Current tobacco use is defined as having used any form of tobacco, on one or more of the past 30 days. Any form of tobacco includes cigarettes, cigars, smokeless or chewing tobacco, pipe tobacco, Bidis (small brown cigarettes from India), or Kreteks, which are more commonly known as "clove cigarettes." Overall prevalence of tobacco use among Bay County middle school students has decreased by 14.7 percent from 19.4 percent in 2000 to 16.5 percent in 2006. Overall prevalence tobacco use among Bay County high school students has decreased by 29.3 percent from 38.3 percent in 2000 to 27.1 percent in 2006.



*Data are available only where point estimates are shown.


Current Cigarette Use

Current cigarette use is defined as having smoked a cigarette on one or more of the past 30 days. The prevalence of cigarette use among Bay County middle school students has decreased by 34.6 percent from 14.4 percent in 2000 to 9.4 percent in 2006. The prevalence of cigarette use among Bay County high school students has decreased by 34.7 percent from 25.3 percent in 2000 to 16.5 percent in 2006.



Current Cigar Use


Current cigar use is defined as having smoked a cigar on one or more of the past 30 days. The prevalence of cigar use among Bay County middle school students has increased by 5.9 percent from 10.1 percent in 2000 to 10.7 percent in 2006. The prevalence of cigar use among Bay County high school students has decreased by 24.4 percent from 22.1 percent in 2000 to 16.7 percent in 2006.



*Data are available only where point estimates are shown.


Current Smokeless Tobacco Use

Current smokeless tobacco use is defined as having used smokeless tobacco on one or more of the past 30 days. The prevalence of smokeless tobacco use among Bay County middle school students has increased 14.5 percent from 5.5 percent in 2000 to 6.3 percent in 2006. The prevalence of smokeless tobacco use among Bay County high school students did not change significantly from 2000 (8.7 percent) to 2006 (8.7 percent).



“Committed Never Smokers”

“Committed never smokers” are students who have never smoked a cigarette and state they will definitely not smoke a cigarette in the future or if a friend offered one. The prevalence of “committed never smokers” in Bay County middle schools has increased by 20.9 percent from 48.7 percent in 2000 to 68.6 percent in 2006. The prevalence of “committed never smokers” in Bay County high schools has increased by 49.8 percent from 33.3 percent in 2000 to 83.1 percent in 2006.




*Data are available only where point estimates are shown.

Tobacco Use Prevention Education (TUPE)

To assess the amount of tobacco use prevention education (TUPE) students receive, survey respondents were asked about what students have learned regarding tobacco in school in the past year. Five questions related to TUPE were asked. Students were asked if they had learned: 1) about tobacco use; 2) about reasons why people of their age smoke; 3) that most people of the same age do not smoke; 4) ways to say “no”; and 5) about the negative effects of smoking, such as yellowed teeth, wrinkles, or bad smell.


Comprehensive TUPE

Students who respond they have been taught about all five of the above-mentioned topics during the past school year are considered having received comprehensive TUPE. Middle school students in Bay County showed a 56.8 percent increase of receiving comprehensive TUPE from 8.1 percent in 2000 to 12.7 percent in 2006. The prevalence of receiving comprehensive TUPE among Bay County high school students has remained stable around 5.7 percent from 2002 to 2006.



Any TUPE

Students who respond that they have been taught about any one of the above-mentioned topics are considered having received “Any TUPE.” Middle school students in Bay County show an 8.6 percent increase of receiving “any TUPE” from 61.4 percent in 2000 to 66.7 percent in 2006. High school students in Bay County show a 30.7 percent increase of receiving “any TUPE” from 39.8 percent in 2000 to 52 percent in 2006.




*Data are available only where point estimates are shown.

Secondhand Smoke Exposure

Secondhand smoke (SHS), also known as environmental tobacco smoke, refers to a combination of side-stream smoke (smoke that is emitted from the end of a lit cigarette, pipe, or cigar) and main-stream smoke (smoke exhaled by a smoker). SHS is the third leading cause of preventable death in the United States.¹ In 2003, Florida passed the Florida Clean Indoor Air Act, which bans smoking in workplaces to try to reduce the number of Floridians negatively impacted by the effects of SHS.

Exposure to Secondhand Smoke


The prevalence of exposure to SHS among Bay County middle school students decreased by 5.1 percent from 67.1 percent in 2000 to 63.7 percent in 2006. The prevalence of exposure to SHS among Bay County high school students has decreased by 14.0 percent from 79.3 percent in 2000 to 68.2 percent in 2006.




Source of Secondhand Smoke Exposure: 2006 Survey Data

The two charts below show the percentage of people who were reported as the source of SHS exposure by Bay County middle and high school students. Middle school students were most likely to be exposed to SHS by their parents, while high school students were equally likely to be exposed to SHS by their parents and their friends.

Bay County Middle School Students



Bay County High School Students




¹ "Secondhand Smoke, Who is at Risk?" Florida Department of Health, Division of Environmental Health, 2006.

*Data are available only where point estimates are shown.


Location of Secondhand Smoke Exposure: 2006 Data

The two charts below show the locations in which Bay County middle and high school students were exposed to SHS. Both middle and high school students were most likely to be exposed to SHS at home.

Bay County Middle School Students



Bay County High School Students




Tobacco Sources

Source of Cigarettes for Minors

Students are asked how they usually acquire their cigarettes. This analysis has been limited to students who are minors less than 18 years of age. The chart below shows how Bay County minors (in either middle or high school) acquire cigarettes. Bay County minors are most likely to get their cigarettes by giving someone else money to purchase them.

Cigarette Sources for Minors




Anti-Tobacco Activism

Participation in an Anti-Tobacco Community Event

The prevalence of Bay County middle school students' participation in anti-tobacco community events has doubled from 15.2 percent in 2000 to 30.5 percent in 2006. The prevalence of Bay County high school students' participation in anti-tobacco community events has increased by 81.3 percent from 8.0 percent in 2000 to 14.5 percent in 2006.


Participation in Community Anti-Tobacco Events*



Students Working Against Tobacco (S.W.A.T.)

The prevalence of Bay County middle school students who served as active members of S.W.A.T. has decreased by 57.0 percent from 20.3 percent in 2000 to 8.7 percent in 2006. The prevalence of Bay County high school students who served as active members of S.W.A.T. has increased 25.9 percent from 5.4 percent in 2000 to 6.8 percent in 2006.

Active Members of SWAT*




*Data are available only where point estimates are shown.

Social Attitudes


Perception that Smokers have More Friends

Students are asked whether or not they believe that young people who smoke have more friends. In 2006, 31.2 percent of Bay County middle school students and 30.3 percent of Bay county high school students definitely did not think that smokers have more friends.



Perception that Smokers Look Cool or Fit In

Students are asked whether or not they believe that smoking helps young people “look cool” or “fit in.” In 2006, 70.9 percent of Bay County middle school students and 73 percent of Bay County high school students definitely did not think that smoking helps young people “fit in” or “look cool.”




*Data are available only where point estimates are shown.

Bay County: 2006 Data at a Glance

Indicator	Middle School				High School			
	Bay County %	95 % CI	State %	95 % CI	Bay County %	95 % CI	State %	95 % CI
Smoked cigarettes on one or more of the past 30 days	9.4	(7.1 - 11.8)	6.6	(6.1 - 7.1)	16.5	(13.6 - 19.4)	15.5	(14.7 - 16.3)
Smoked cigars on one or more of the past 30 days	10.7	(8.2 - 13.2)	6.3	(5.9 - 6.8)	16.7	(13.8 - 19.7)	14.0	(13.3 - 14.7)
Used smokeless tobacco one or more of the past 30 days	6.3	(4.3 - 8.3)	3.4	(3.1 - 3.8)	8.7	(6.5 - 11.0)	6.0	(5.5 - 6.6)
Used any form of tobacco on one or more of the past 30 days	16.5	(4.1 - 7.8)	11.3	(10.6 - 12.0)	27.1	(13.5 - 19.5)	23.9	(22.9 - 25.0)
Have never smoked a cigarette and will definitely not smoke a cigarette in the future or if a friend offered one	58.9	(54.8 - 63.0)	60.9	(59.8 - 61.9)	49.9	(45.8 - 53.9)	48.6	(47.5 - 49.8)
Exposed to secondhand smoke during the past 7 days	63.7	(59.8 - 67.6)	56.3	(55.1 - 57.6)	68.2	(64.5 - 71.8)	64.1	(63.0 - 65.3)
Received Comprehensive tobacco use prevention education (TUPE)	12.7	(10.0 - 15.3)	16.7	(15.6 - 17.8)	5.6	(3.8 - 7.3)	8.8	(8.2 - 9.4)
Received any tobacco use prevention education (TUPE)	66.7	(62.9 - 70.5)	71.3	(69.6 - 72.9)	52.0	(48.1 - 55.8)	53.3	(51.9 - 54.6)
Participated in a community event	30.5	(26.9 - 34.2)	23.8	(23.0 - 24.6)	14.5	(11.8 - 17.2)	14.1	(13.5 - 14.7)
Served as an active member of SWAT	8.7	(6.5 - 11.0)	4.0	(3.6 - 4.4)	6.8	(4.9 - 8.7)	3.4	(3.1 - 3.7)
<u>Definitely did not think that smokers have more friends</u>	31.2	(27.5 - 34.9)	32.6	(31.8 - 33.4)	30.3	(26.7 - 33.9)	32.0	(31.2 - 32.9)
<u>Definitely did not think that smoking helps young people "look cool" or "fit in"</u>	70.9	(67.3 - 74.6)	72.6	(71.9 - 73.4)	73.0	(69.6 - 76.5)	71.1	(70.3 - 71.9)

2006 Bay County Florida Youth Tobacco Survey Report



The collection and reporting of county-level survey data would not have been possible without the collaboration and tireless efforts of many. We thank our colleagues at: Florida Department of Health; Florida Department of Children and Families; Florida Department of Education; Florida Department of Juvenile Justice; Florida's Executive Office of the Governor, Office of Drug Control; The Centers for Disease Control and Prevention, Office on Smoking and Health; and ORC Macro. Special thanks go to our friends in the County Healthy Departments and local School District Offices. Most especially we thank the principals, teachers, and students who participated in the survey and made this information possible.

For more information about the FYTS, please contact: Ms. Jamie Weitz, M.S., Florida Youth Survey Coordinator, at (850) 245-4444, extension 2424 or by e-mail at Jamie_Weitz@doh.state.fl.us. Also, visit our website at: http://www.doh.state.fl.us/disease_ctrl/epi/Chronic_Disease/FYTS/Intro.htm .

